

Supplementary Table S1. Oligonucleotides used for qPCR and RNAi

small RNA RT-qPCR	
Neg_ctrl_RT	CTAGCCTTGAGTCGAGCAGGTCTAGACTACTTACT
Neg_ctrl_FWD	AA+TGTATA+CATTGTC+AG
Neg_ctrl_REV	GTCGAGCAGGTCTAGACTACTTACT
miR-AS1_RT	CTGGTGTCTGGAGTCGGCAATGGTGCCAC
miR-AS1_FWD	G+AA+GAG+CAGCAGAAG
miR-AS1_REV	CTGGTGTCTGGAGTCGGCAA
let7a-RT	TGATCGGTAGTAGGACCATCAACTATAC
let7a-FWD	T+GA+GGTAGTAGGTTG
let7a-REV	TGATCGGTAGTAGGACCATC
mRNA RT-qPCR	
RdRp_FWD	GTGAAATGGTCATGTGTGGCGG
RdRp_REV	CAGATGTTAAACACACTATTAGCATA
GAPDH_FWD	GAAATCCCATCACCATCTTCCAG
GAPDH_REV	GAGCCCCAGCCTTCTCCATG
siRNAs	
Ambion	Silencer Negative control #AM4611
SARS_CoV-2-miR-1	GAAGAGCAGCAGAAGUGGCACCA
SARS_CoV-2-miR-1_as	GUGCCACUUCUGCUGCUCUUCUU

Bases prepended by a "+" are LNA modified