

Heterobimetallic One-Dimensional Coordination Polymers $M^I Cu^{II}$ (M = Li and K) Based on Ferromagnetically Coupled Di- and Tetracopper(II) Metallacyclophanes

Tamyris T. da Cunha¹, Willian X. C. Oliveira¹, Emerson F. Pedroso², Francesc Lloret³, Miguel Julve³, Cynthia L. M. Pereira^{1,*}

¹ Departamento de Química, Instituto de Ciências Exatas, Universidade Federal de Minas Gerais. Av. Antônio Carlos 6627, Pampulha, Belo Horizonte, Minas Gerais, 31270-901 Brazil; tamyris.cunha@gmail.com (T.T.C.); williancoliveira@yahoo.com.br (W.X.C.O.).

² Centro Federal de Educação Tecnológica de Minas Gerais, Av. Amazonas 5855, 30421169, Belo Horizonte, Minas Gerais; pedroso@cefetmg.br (E.F.P)

³ Departamento de Química Inorgánica/Instituto de Ciencia Molecular (ICMol), Universitat de València, C/ Catedrático José Beltrán 2, 46980 Paterna (València), Spain; Francisco.Lloret (F.L); Miguel.Julve@uv.es (M.J.)

* Corresponding author: cynthialopes@ufmg.br (C.L.M.P.), Tel.: +55-31-3409-7556




Figure S1. TG curve of 1.




Figure S2. TG curve of 2.




Figure S3. Experimental and Simulated X-ray powder diffraction patterns for **1**.




Figure S4. Experimental and simulated X-ray powder diffraction patterns for **2**.




Figure S5. Perspective view of the dicopper [3,3] metallacyclophane-type fragment of **1** with atomic displacement ellipsoids at 50% of probability. Circles represents isotropically treated disordered atoms. Color code: carbon (grey), nitrogen (sky blue), lithium (purple) and copper (green).




Figure S6. Representation of a fragment of the structure of **2** with atomic displacement ellipsoids at 50% of probability. Circles represents isotropically treated disordered atoms. Color code: carbon (grey), nitrogen (sky blue), lithium (purple) and copper (green) [Symmetry code: (i) = $5/2-x, y, 2-z$].